

Climate change could expose volcanic regions, increase eruptions: Study

---

27/11/2017



Climate change caused by humans is rapidly melting ice in volcanically active regions, which could lead to increased volcano eruptions, a study has found.

Climate change caused by humans is rapidly melting ice in volcanically active regions, which could lead to increased volcano eruptions, a study has found. The study, led by researchers at the University of Leeds in the UK, found that there was less volcanic activity in Iceland when glacier cover was more extensive and as the glaciers melted volcanic eruptions increased due to subsequent changes in surface pressure.

“Climate change caused by humans is creating rapid ice melt in volcanically active regions. In Iceland, this has put us on a path to more frequent volcanic eruptions,” said Graeme Swindles from the School of Geography at Leeds. The study examined Icelandic volcanic ash preserved in peat deposits and lake sediments and identified a period of significantly reduced volcanic activity between 5,500 and 4,500 years ago.

This period came after a major decrease in global temperature, which caused glacier growth in Iceland. The findings, published in the journal *Geology*, found

there was a time lag of roughly 600 years between the climate event and a noticeable decrease in the number of volcanic eruptions. The study suggests that perhaps a similar time lag can be expected following the more recent shift to warmer temperatures.



Icelandic volcanism is controlled by complex interactions between rifts in continental plate boundaries, underground gas and magma build-up and pressure on the volcano's surface from glaciers and ice. (File Photo)

Iceland's volcanic system is in process of recovering from the 'Little Ice Age' – a recorded period of colder climate roughly between the years 1500 and 1850 AD, researchers said. Since the end of the Little Ice Age, a combination of natural and human caused climate warming is causing Icelandic glaciers to melt again, they said. "The human effect on global warming makes it difficult to predict how long the time lag will be but the trends of the past show us more eruptions in Iceland can be expected in the future," Swindles said.

"These long term consequences of human effect on the climate is why summits like COP are so important. It is vital to understand how actions today can impact future generations in ways that have not been fully realised, such as more ash clouds over Europe, more particles in the atmosphere and problems for aviation," he said. Icelandic volcanism is controlled by complex interactions between rifts in continental plate boundaries, underground gas and magma build-up and pressure on the volcano's surface from glaciers and ice.

Changes in surface pressure can alter the stress on shallow chambers where magma builds up. "When glaciers retreat there is less pressure on the Earth's

surface. This can increase the amount of mantle melt as well as affect magma flow and how much magma the crust can hold,” said Ivan Savov, from the School of Earth & Environment at Leeds. “Even small changes in surface pressure can alter the likelihood of eruptions at ice-covered volcanos,” said Savov.

---