

Culex mosquitoes do not transmit zika virus, study finds

04/10/2016



A Biosecurity Research Institute study has found important results in the fight against Zika virus: *Culex* mosquitoes do not appear to transmit Zika virus.

Researchers at Kansas State University's Biosecurity Research Institute studied *Culex* species mosquitoes from across the country, including Vero Beach in Florida, which is near Miami-Dade County where mosquitoes are spreading Zika virus.

The research, "*Culex* species mosquitoes and Zika virus," appears in the journal *Vector-Borne and Zoonotic Diseases* and involves researchers from Rutgers University, the University of Florida and the U.S. Department of Agriculture.

The findings are important for controlling Zika virus in Florida and preventing its spread to other parts of the country, said Dana Vanlandingham, lead author and assistant professor of virology in the College of Veterinary Medicine.

"It's very important to know that *Culex* mosquitoes are not able to transmit Zika," Vanlandingham said. "It enables people to target their control strategies so that they aren't wasting time and effort on a mosquito that isn't transmitting Zika virus."

It is the first Zika virus research publication from the Biosecurity Research Institute. Before this study, *Culex* mosquitoes' role in Zika virus was unclear. By studying *Culex* mosquitoes over a period of time, the researchers found that Zika virus did not multiply and instead disappeared in the species.

"This is great news," said Stephen Higgs, co-author and director of the Biosecurity Research Institute. "We can check this particular group of mosquitoes off the list here in the U.S. and focus efforts of control on the mosquitoes that we know can infect, like *Aedes aegypti* and *Aedes albopictus*."

The Centers for Disease Control and Prevention has identified *Aedes aegypti*, or yellow fever mosquito, and *Aedes albopictus*, or Asian tiger mosquito, as two species that transmit Zika virus. Both mosquitoes are widely distributed in the U.S. and are present in Kansas.

*Culex* mosquitoes are brown mosquitoes, while *Aedes aegypti* are black and *Aedes albopictus* are black and white. *Culex* mosquitoes transmit West Nile virus and Japanese encephalitis and live outside. *Aedes aegypti* and *Aedes albopictus* can live in and around houses in plant trays, spare containers or gutters.

"We need to know which mosquitoes to target and which mosquitoes not to target because mosquitoes live in different environments," said Vanlandingham, whose research focuses on zoonotic viruses -- such as Japanese encephalitis and chikungunya. "Some mosquitoes are found outside and some are more in people's homes. You need to know this in order to target your efforts."

Both Vanlandingham and Higgs emphasize the importance of personal responsibility in stopping the spread of Zika virus. Homeowners can get rid of small pools of water where mosquitoes breed and should use mosquito repellent as personal protection.

While a startup fund from the university's College of Veterinary Medicine provided funding for this Biosecurity Research Institute study, there is still a need for additional national funding to support research that stops Zika virus, said Higgs, who also has studied chikungunya, a mosquito-borne virus that has a similar transmission cycle to that of Zika virus.

"We thought that this research is so important with what is going on that we were able to use startup funding," Higgs said. "This research is basic research because we don't know some of the most fundamental information about mosquitoes. Applied research -- such as vaccines and diagnostics -- are obviously very important, but there is a need for funding basic research as well."